

Table 1

Compaision of patient's data between normal patients and alcoholic liver cirrhosis patients devided by Child-Pugh classification.

Variants	All patients (N=97)	Normal patients (N=19)	Child-Pugh grade A (N=56)	Child-Pugh grade B (N=15)	Child-Pugh grade C (N=7)	P value
Age (Year)	67.08±11.75	67.41±9.16	68.95±12.47	63.20±9.80	59.71±12.47	0.12 ¹⁾
Sex (M/F)	79/16	12/5	49/7	13/2	5/2	0.26 ²⁾
Child-Pugh score	-	-	5.36±0.48	7.87±0.92	11.43± 1.62	-
AST (IU/L)	34(24-52.5)	21(18-24)	36(25-51)	48(37-83.5)	57(55.5-155)	<0.001 ³⁾
ALT (IU/L)	22(15-36)	12(6-16)	22.5(15.5-35)	32(21-41.5)	36(22-55)	0.0011 ³⁾
γ-GTP (IU/L)	102.43±65.47	23.19±12.98	115.71±73.21	128.00±81.87	167.32±125. 11	0.032 ¹⁾
PT (%)	83.83±21.97	100.24±11.49	88.25±18.01	62.47±17.35	54.46± 21.09	<0.001 ¹⁾
Albumin (g/dL)	3.87±0.70	4.44±0.24	4.01±0.52	3.34±0.59	2.51± 0.41	<0.001 ¹⁾
T-Bil (mg/dL)	1.67±1.31	0.85±0.34	1.25±0.55	2.35±1.23	6.06±3.51	<0.001 ¹⁾

AST; aspartate aminotransferase, ALT; alanine aminotransferase, γ-GTP; γ-glutamyl transpeptidase, PT; prothrombin time, T-Bil; total bilirubin.

- 1) Comparisons between groups were made using one-way analysis of variance (ANOVA).
- 2) Comparisons between groups were made using Fisher's exact analysis.
- 3) Comparisons between groups were made using Kruskal-Wallis test.